

Finiteness properties of automorphism groups of right-angled Artin groups

Ruth Charney and Karen Vogtmann

ABSTRACT

We study the algebraic structure of the outer automorphism group of a general right-angled Artin group. We show that this group is virtually torsion-free and has finite virtual cohomological dimension. This generalizes results proved in [?] for two-dimensional right-angled Artin groups.

1. Introduction

Associated to a finite simplicial graph Γ is a right-angled Artin group A_Γ whose generators are the vertices of Γ and whose relations are commutators between adjacent vertices. These groups range from free groups (when Γ has no edges) to free abelian groups (when Γ is a complete graph), hence their outer automorphism groups interpolate between $Out(F_n)$ and $GL_n(\mathbb{Z})$. The automorphism groups of free groups and free abelian groups have many properties in common. In particular, they are virtually torsion-free, have finite virtual cohomological dimension, and satisfy a Tits alternative (every subgroup is either virtually solvable or contains a free group of rank 2). One is naturally led to ask whether the same is true for automorphism groups of right-angled Artin groups.

In [?] we began a study of the groups $Out(A_\Gamma)$ by analyzing the case when the defining graph Γ is connected and has no triangles. We used both algebraic and geometric methods to establish cohomological finiteness results as well as to prove that the Tits alternative holds. The key algebraic tools we used were certain projection homomorphisms. To define these homomorphisms, we introduced a partial ordering and an equivalence relation on the vertices of Γ . For each maximal vertex v , the projection homomorphism P_v was defined on a finite index subgroup of $Out(A_\Gamma)$ and took its image in $Out(A_{lk(v)})$ where $lk(v)$ denotes the link of v in Γ . These projection homomorphisms were assembled into a single homomorphism P , and the kernel of P was computed precisely.

In this paper we show how similar projection homomorphisms can be defined for general right-angled Artin groups. We then determine enough information about the kernel of the composite homomorphism to be able to construct an inductive proof that $Out(A_\Gamma)$ is virtually torsion-free. Thus the virtual cohomological dimension of $Out(A_\Gamma)$ is defined, and we go on to prove that it is finite for any finite simplicial graph Γ .

2000 *Mathematics Subject Classification* 20F36 (primary).

R. Charney was partially supported by NSF grant DMS 0705396. K. Vogtmann was partially supported by NSF grant DMS 0705960

2. Join subgroups

Let Γ be a connected, simplicial graph with vertex set V , and A_Γ the associated right-angled Artin group. For two vertices $v, w \in V$, let $d(v, w)$ denote the distance from v to w in Γ . The link $lk(v)$ is the subgraph spanned by vertices at distance one from v , and the star $st(v)$ is the subgraph spanned by vertices at distance at most one from v . Note that $v \in lk(w)$ if and only if $w \in lk(v)$, and $v \in st(w)$ if and only if $w \in st(v)$.

We first need to define an appropriate partial ordering and equivalence relation on vertices in Γ , and prove some of their basic properties. For $v, w \in V$, set $v \leq w$ if $lk(v) \subseteq st(w)$. Note that this may occur in three ways:

- (1) $v = w$,
- (2) $d(v, w) = 1$ and $st(v) \subseteq st(w)$,
- (3) $d(v, w) = 2$ and $lk(v) \subseteq lk(w)$.

LEMMA 2.1. *If $u \leq v \leq w$ and $d(u, v) = 1$, then $d(v, w) \leq 1$.*

Proof. Since $d(u, v) = 1$, $u \in lk(v) \subseteq st(w)$. Therefore $w \in st(u)$, i.e. either $w = u$ or $w \in lk(u) \subseteq st(v)$. In either case $d(v, w) \leq 1$. \square

LEMMA 2.2. *The relation \leq is transitive on the vertices of Γ .*

Proof. Suppose $u \leq v \leq w$. We need to check that $u \leq w$. If $d(u, v) = 0$ or 2 , then $lk(u) \subseteq lk(v) \subseteq st(w)$, so this is immediate. If $d(u, v) = 1$, then by Lemma ??, $d(v, w) \leq 1$. Since we already know that $lk(v) \subseteq st(w)$, this gives $st(v) \subseteq st(w)$, so $lk(u) \subseteq st(v) \subseteq st(w)$. \square

Now define a relation on vertices by $v \sim w$ if $v \leq w$ and $w \leq v$, or equivalently, if one of the following holds,

- (1) $v = w$,
- (2) $d(v, w) = 1$ and $st(v) = st(w)$,
- (3) $d(v, w) = 2$ and $lk(v) = lk(w)$.

It follows from Lemma ?? that this is an equivalence relation and that \leq induces a partial ordering on the set of equivalence classes. We denote the equivalence class of v by $[v]$.

LEMMA 2.3. *All elements of $[v]$ are the same distance from each other.*

Proof. If $x, y \in [v]$ and $d(x, y) = 1$, then any other $z \in [v]$ satisfies $y \leq x \leq z$, so by Lemma ??, $d(x, z) = 1$. \square

Thus $[v]$ generates either a free abelian subgroup of A_Γ or a non-abelian free subgroup. Let V_{ab} denote the set of vertices v with $A_{[v]}$ abelian, and V_{fr} the set of vertices with $A_{[v]}$ non-abelian.

LEMMA 2.4. *Suppose $[v_1] < [v_2] < \dots < [v_k]$. Then there exists j , $0 \leq j \leq k$, such that $d(v_i, v_{i+1}) = 2$ for all $i < j$ and $d(v_i, v_{i+1}) = 1$ for all $i \geq j$. Moreover, for $i > j$, $v_i \in V_{ab}$.*

Proof. This is immediate from Lemma ??.

□

It is not necessarily the case that v_i is non-abelian for $i \leq j$, since an abelian $[v]$ can be less than a non-abelian $[w]$ if $[v]$ consists of just a single vertex.

The projection homomorphisms in [?] mapped first to groups associated to certain *maximal join subgraphs* containing a given maximal vertex, then to groups associated to the link of the vertex. In the general case, the definitions of these subgraphs must be modified as follows.

For two simplicial graphs Θ_1 and Θ_2 , let $\Theta_1 * \Theta_2$ denote their join, that is, the graph formed by joining every vertex of Θ_1 to every vertex of Θ_2 by an edge. Note that if $\Theta = \Theta_1 * \Theta_2$, then $A_\Theta = A_{\Theta_1} \times A_{\Theta_2}$.

For a vertex v in Γ , define

$$\begin{aligned} L_{[v]} &= lk(v) \setminus [v] \\ J_{[v]} &= L_{[v]} * [v] \end{aligned}$$

$J_{[v]}$ is called the *join associated to* $[v]$. Note that it is always the case that $st(v) \subseteq J_{[v]}$ and equality holds if and only if $v \in V_{ab}$. Note also, that the relation $w \in L_{[v]}$ is symmetric; that is, w lies in $L_{[v]}$ if and only if v lies in $L_{[w]}$.

The special subgroup generated by $J_{[v]}$ is a direct product

$$A_{J_{[v]}} = A_{L_{[v]}} \times A_{[v]}$$

where the second factor is either free or free abelian.

In order to understand the relation between various equivalence classes of vertices and their associated joins, it is useful to consider a graph Γ_0 defined as follows. The vertices of Γ_0 are the equivalence classes $[v]$ which are maximal with respect to the partial order \leq . Two vertices $[v]$ and $[w]$ are joined by an edge in Γ_0 if and only if v and w are adjacent in Γ . It follows from the definition of the equivalence relation that adjacency is independent of choice of representative. We remark that a similar graph, based on this same equivalence relation, is described in recent work of Duncan, Kazachkov, and Remeslennikov [?].

LEMMA 2.5. Γ_0 is a connected graph and every $w \in V$ lies in the join $J_{[v]}$ associated to some $[v]$ in Γ_0 .

Proof. To prove connectivity of Γ_0 , let $[v]$ and $[v']$ be vertices of Γ_0 . Let $v = v_0, v_1, \dots, v_n = v'$ be an edgepath in Γ . We proceed by induction on n . If $n = 1$, then $[v], [v']$ are adjacent in Γ_0 . Suppose $n > 1$. Choose $[w]$ such that $[w]$ is maximal and $[v_1] \leq [w]$. Then v and v_2 lie in $lk(v_1) \subset st(w)$. So either $[v] = [w]$ or $[v]$ is adjacent to $[w]$ in Γ_0 , and there is an edgepath w, v_2, \dots, v_n in Γ . By induction, $[w]$ is connected to $[v']$ by a path in Γ_0 .

For the second statement of the lemma, note that if w is in the link of u , then for any v with $[u] \leq [v]$, $w \in lk(u) \subset st(v) \subset J_{[v]}$. □

For a subgraph Θ of Γ , denote the normalizer, centralizer and center of A_Θ respectively by $N(\Theta)$, $C(\Theta)$ and $Z(\Theta)$. It follows from a theorem of E. Godelle [?] (see also [?], Proposition 2.2), that

$$N(\Theta) = A_{\Theta \cup \Theta^\perp} \quad C(\Theta) = A_{\Theta^\perp} \quad Z(\Theta) = A_{\Theta \cap \Theta^\perp}.$$

where Θ^\perp is the set of vertices commuting with all elements of Θ . We will be particularly interested in the case of $\Theta = J_{[v]}$.

LEMMA 2.6. *If $[v]$ is maximal, then the centralizer, center, and normalizer of $A_{J_{[v]}}$ are given by*

$$C(J_{[v]}) = Z(J_{[v]}) = A_{[v]} \text{ if } v \in V_{ab} \text{ and } \{1\} \text{ if } v \in V_{fr}$$

$$N(J_{[v]}) = A_{J_{[v]}}.$$

Proof. Suppose $[v]$ is maximal and some u commutes with all of $J_{[v]}$. Then $lk(v) \subseteq J_{[v]} \subseteq st(u)$, so by maximality of $[v]$, $[u] = [v]$. Thus J_v^\perp is contained in $[v]$ and the lemma follows from the descriptions above. \square

LEMMA 2.7. *Suppose $[v]$ and $[w]$ are adjacent vertices in Γ_0 . Let $J_{v,w}$ denote the intersection of $J_{[v]}$ and $J_{[w]}$. Then*

$$C(J_{v,w}) = Z(J_{v,w}) = Z([v]) \times Z([w]) \times Z(L_{[v]} \cap L_{[w]})$$

$$N(J_{v,w}) = A_{J_{v,w}}.$$

Proof. Since $[v]$ and $[w]$ are adjacent, $[v] \subset L_{[w]}$ and $[w] \subset L_{[v]}$. Thus, $J_{v,w}$ decomposes as a join, $J_{v,w} = [v] * [w] * (L_{[v]} \cap L_{[w]})$. Any generator commuting with both $[v]$ and $[w]$ lies in $J_{v,w}$, so $J_{v,w}^\perp \subset J_{v,w}$. The lemma now follows from the formulas above. \square

3. Projection homomorphisms

In this section we describe the projection homomorphism P , which will be defined on a finite-index subgroup of $Out(A_\Gamma)$. We begin by reviewing the work of M. Laurence [?]. Building on the work of H. Servatius [?], Laurence described a finite set of generators for $Aut(A_\Gamma)$ as follows.

- (1) *Inner automorphisms* conjugate the entire group by some generator v .
- (2) *Symmetries* are induced by symmetries of Γ and permute the generators.
- (3) *Inversions* send a standard generator of A_Γ to its inverse.
- (4) *Transvections* occur whenever $v \leq w$, and send $v \mapsto vw$.
- (5) *Partial conjugations* occur whenever removing the (closed) star of a vertex v disconnects Γ . If this happens, a partial conjugation conjugates all of the generators in one component of $\Gamma - st(v)$ by v .

DEFINITION 3.1. The subgroup of $Aut(A_\Gamma)$ generated by inner automorphisms, inversions, partial conjugations and transvections is called the *pure automorphism group* and is denoted $Aut^0(A_\Gamma)$. The image of $Aut^0(A_\Gamma)$ in $Out(A_\Gamma)$ is the group of *pure outer automorphisms* and is denoted $Out^0(A_\Gamma)$.

The subgroups $Aut^0(A_\Gamma)$ and $Out^0(A_\Gamma)$ are easily seen to be normal and of finite index in $Aut(A_\Gamma)$ and $Out(A_\Gamma)$ respectively. We remark that if A_Γ is a free group or free abelian group, then $Aut^0(A_\Gamma) = Aut(A_\Gamma)$.

PROPOSITION 3.2. *Let $[v]$ be a maximal equivalence class. Then any $\phi \in Out^0(A_\Gamma)$ has a representative $\phi_v \in Aut^0(A_\Gamma)$ which preserves both $A_{[v]}$ and $A_{J_{[v]}}$.*

Proof. It suffices to check that the proposition holds for each of the generators of $Out^0(A_\Gamma)$. It is clear for inversions. Represent $\hat{\phi}$ by some automorphism $\hat{\phi} \in Aut^0(A_\Gamma)$.

Partial conjugations: Suppose $\hat{\phi}$ is a partial conjugation by a generator w . If w is not in $J_{[v]}$, then $d(w, v) \geq 2$, so $st(w) \cap J_{[v]} \subseteq L_{[v]}$. If $[v]$ is abelian, $J_{[v]} - st(w)$ is clearly connected. If $[v]$ is nonabelian, then maximality of $[v]$ implies that $st(w) \cap J_{[v]}$ is not all of $L_{[v]}$, so that in this case too $J_{[v]} - st(w)$ is connected. Thus $\hat{\phi}$ is either trivial on $A_{J_{[v]}}$ or acts as conjugation by w on all of $A_{J_{[v]}}$. Composing $\hat{\phi}$ with the inner automorphism associated to w produces the desired ϕ_v .

If $w \in J_{[v]} = [v] * L_{[v]}$, then conjugation by w preserves the two factors $A_{[v]}$ and $A_{L_{[v]}}$, hence the same is true for the partial conjugation $\hat{\phi}$, so we may take $\phi_v = \hat{\phi}$.

Transvections: Suppose $u \leq w$ and $\hat{\phi}$ is the transvection $u \mapsto uw$. If u is not in $J_{[v]}$ then $\hat{\phi}$ is the identity on $A_{J_{[v]}}$. If $u \in L_{[v]}$, then $\hat{\phi}$ fixes $A_{[v]}$, and $v \in lk(u) \subseteq st(w)$, so $w \in st(v) \subseteq J_{[v]}$ and $\hat{\phi}$ preserves $A_{J_{[v]}}$. If $u \in [v]$, then maximality of $[v]$ implies $[v] = [u] = [w]$, so the transvection preserves both $A_{[v]}$ and $A_{J_{[v]}}$. In all cases we may take $\phi_v = \hat{\phi}$. \square

The representative ϕ_v described in Proposition ?? is well defined up to conjugation by an element of the normalizer of $A_{J_{[v]}}$. In light of Lemma ??, the restriction of ϕ_v to $A_{J_{[v]}}$ is well defined up to an inner automorphism of $A_{J_{[v]}}$. Moreover, since ϕ_v preserves $A_{[v]}$, it projects to an automorphism of $A_{L_{[v]}} \cong A_{J_{[v]}}/A_{[v]}$. This immediately gives the following corollary.

COROLLARY 3.3. *For every maximal $[v]$ there is a restriction homomorphism*

$$R_{[v]} : Out^0(A_\Gamma) \rightarrow Out(A_{J_{[v]}})$$

and a projection homomorphism

$$P_{[v]} : Out^0(A_\Gamma) \rightarrow Out(A_{L_{[v]}})$$

We now assemble these homomorphisms, one for each maximal equivalence class $[v]$, to obtain a restriction homomorphism

$$R = \prod R_{[v]} : Out^0(A_\Gamma) \rightarrow \prod Out(A_{J_{[v]}})$$

and a projection homomorphism

$$P = \prod P_{[v]} : Out^0(A_\Gamma) \rightarrow \prod Out(A_{L_{[v]}}).$$

4. The kernel of the projection homomorphism

In order to use the projection homomorphism P to obtain information about $\text{Out}(A_\Gamma)$ we need to understand basic properties of its kernel. We first consider the kernel of the restriction homomorphism R .

THEOREM 4.1. *The kernel K of the homomorphism R is a finitely generated free abelian group.*

Proof. If Γ_0 consists of a single vertex $[v]$, then $\Gamma = J_{[v]}$ so the kernel is trivial. So assume that there is more than one maximal equivalence class.

By definition, elements of K are outer automorphisms. We begin by choosing a canonical automorphism to represent each element of K . First note that for any element ϕ of K and any maximal $[v]$, we can choose a representative automorphism ϕ_v such that the restriction of ϕ_v to $A_{J_{[v]}}$ is the *identity* map. This representative is unique up to conjugation by an element of the centralizer $C(J_{[v]})$. In particular, if $[v]$ is non-abelian, then ϕ_v is unique.

Suppose $V_{f_r} \neq \emptyset$. Choose a non-abelian class $[y]$. Then for any element ϕ of K , we take $\phi_0 = \phi_y$ as our canonical representative. If $V_{f_r} = \emptyset$, choose a pair of adjacent maximal classes $[y]$ and $[z]$. Note that $\phi_y = c(a)\phi_z$ where $c(a)$ denotes conjugation by an element $a \in C(J_{y,z})$. By Lemma ??, $a = rs$ for some $r \in A_{[y]}$, $s \in A_{L_{[y]}}$. Set

$$\phi_0 = c(r^{-1})\phi_y = c(s)\phi_z.$$

Then ϕ_0 has the property that (i) it restricts to the identity on vertices of $J_{[y]}$ and (ii) it acts on $J_{[z]}$ as conjugation by an element of $A_{L_{[y]}}$. If ϕ_1 is any other representative of ϕ with these two properties, then it differs from ϕ_0 by conjugation by an element of $A_{[y]} \cap A_{L_{[y]}} = \{1\}$. Thus, ϕ_0 is the unique such representative and we designate it as our canonical representative.

The properties which characterize canonical representatives are preserved under composition, thus the map $\phi \mapsto \phi_0$ defines a homomorphism of K into $\text{Aut}(A_\Gamma)$. For the remainder of the proof we view K as a subgroup of the automorphism group by identifying ϕ with ϕ_0 .

To prove the theorem we will define a homomorphism f of K into a free abelian group and prove that f is injective. Recall that V is the set of vertices of Γ so abelianizing gives a homomorphism $A_\Gamma \rightarrow \mathbb{Z}^V$. Denote the abelianization of g by \bar{g} . For each maximal equivalence class $[v]$, define a homomorphism $f_v : K \rightarrow \mathbb{Z}^{V-[v]}$ as follows. An element $\phi \in K$, acts on $J_{[v]}$ as conjugation by some $g \in A_\Gamma$. The element g is unique up to multiplication by an element of $C(J_{[v]}) \subset A_{[v]}$, thus \bar{g} determines a well defined element of $\mathbb{Z}^{V-[v]}$ (which by abuse of notation we will also denote \bar{g}).

We claim that $f_v(\phi) = \bar{g}$ is a homomorphism. For suppose ρ is another element of K and suppose ρ acts on $J_{[v]}$ as conjugation by h . Then $\phi \circ \rho$ acts on $J_{[v]}$ as conjugation by $\phi(h)g$. Since ϕ takes every generator to a conjugate of itself, it leaves the abelianization unchanged. That is, $\overline{\phi(h)} = \bar{h}$, so $f_v(\phi \circ \rho) = \overline{\phi(h)g} = \bar{g} + \bar{h} = f_v(\phi) + f_v(\rho)$.

Now consider the product homomorphism $f = \prod f_v$ taken over all maximal equivalence classes $[v]$,

$$f : K \rightarrow \prod \mathbb{Z}^{V-[v]}.$$

To complete the proof, we will show that f is injective. Suppose ϕ lies in the kernel of f and suppose $[v]$ and $[w]$ are adjacent maximal classes. If ϕ acts on $J_{[v]}$ as conjugation by g_v and on $J_{[w]}$ as conjugation by g_w , then $g_w^{-1}g_v$ lies in the centralizer of $J_{v,w}$, namely in the abelian group $Z([v]) \times Z([w]) \times Z(L_{[v]} \cap L_{[w]})$. Since $f_v(\phi) = \bar{g}_v = 0$ and $f_w(\phi) = \bar{g}_w = 0$, the exponent sum of

any $u \in L_{[v]} \cap L_{[w]}$ is zero in both g_v and g_w , hence also in $g_w^{-1}g_v$. It follows that $g_w^{-1}g_v$ lies in $Z([v]) \times Z([w])$.

Now consider an edge path in Γ_0 from the base vertex $[y]$ to an arbitrary vertex $[v]$,

$$[y] = [v_0], [v_1], \dots, [v_n] = [v],$$

and suppose ϕ acts on $J_{[v_i]}$ as conjugation by $g_i \in A_\Gamma$. Our canonical representative ϕ was chosen so that $g_0 = 1$. By the discussion above, $g_n = (g_0^{-1}g_1)(g_1^{-1}g_2) \dots (g_{n-1}^{-1}g_n)$ is of the form $g_n = a_0a_1a_2 \dots a_n$, where $a_i \in Z([v_i])$. Since ϕ lies in the kernel of f , $f_{[v]}(\phi) = 0$, so all of the a_i 's, except possibly the last one a_n , are trivial. Thus, ϕ acts on $J_{[v]}$ as conjugation by an element of its center, $Z([v])$, i.e., ϕ acts trivially on $J_{[v]}$. Since $[v]$ was arbitrary, we conclude that ϕ is the identity automorphism. \square

In the case that Γ is connected and has no triangles, a stronger version of this theorem is proved in the authors' previous paper with J. Crisp [?]. In that case, we give the exact rank of K and determine an explicit set of partial conjugations which generate K .

Now let K_P be the kernel of the projection homomorphism

$$P = \prod P_{[v]} : \text{Out}^0(A_\Gamma) \rightarrow \prod \text{Out}(A_{L_{[v]}}).$$

Define a vertex v to be *leaf-like* if

- (1) $L_{[v]}$ contains a unique maximal class $[w]$, and
- (2) $[v] < [w]$.

(An easy exercise shows that if Γ has no triangles, then a vertex is leaf-like if and only if it is a leaf, i.e. has valence 1.) Since $d(v, w) = 1$, it follows from Lemma ?? that w must belong to V_{ab} . Since $[v] < [w]$, there is a transvection $t(v, w)$ taking $v \mapsto vw$. We will call this a *leaf transvection*.

THEOREM 4.2. *Assume Γ_0 has at least two vertices. Then the kernel K_P is a free abelian group generated by K and the set of leaf transvections.*

Proof. From the definition of “leaf-like”, it is easy to see that the leaf transvections $t(v, w)$ generate a free abelian group contained in the kernel of K_P and that this subgroup has trivial intersection with K . Thus, it suffices to show that leaf transvections commute with K and that K_P is generated by K and the leaf transvections.

Since every element of K sends each generator to a conjugate of itself, Theorem 2.2 of [?] says that K lies in the subgroup generated by partial conjugations. We claim that $t(v, w)$ commutes with all partial conjugations and hence with all of K . The only case in which this could fail is the case of a partial conjugation by u where v and w lie in different components of $\Gamma \setminus st(u)$. But this is impossible since v and w are connected by an edge. If the edge lies in $st(u)$, then so do v and w , so u commutes with both of them.

It remains to show that K and the leaf transvections generate all of K_P . Suppose $\phi \in K_P$. Let $[w]$ be maximal and ϕ_w be a representative automorphism preserving $A_{[w]}$ and $A_{J_{[w]}}$. Since inner automorphisms by elements of $L_{[w]}$ also preserve these subgroups, we may assume that ϕ_w projects to the identity automorphism on $A_{L_{[w]}}$, so that for any $v \in L_{[w]}$, $\phi_w(v) = vg$ for some $g \in A_{[w]}$. If $[u]$ is another maximal equivalence class with $v \in L_{[u]}$, then we also have $\phi_u(v) = vh$ for some $h \in A_{[u]}$. However, ϕ_u and ϕ_v differ by an inner automorphism. Since no (non-trivial) element of $A_{[w]}$ is conjugate to an element of $A_{[u]}$, this is impossible unless $g = h = 1$.

So suppose that $g \neq 1$ and $[w]$ is the unique maximal equivalence class with $v \in L_{[w]}$ (or equivalently, $[w]$ is the unique maximal class in $L_{[v]}$). We claim that $C(v) \subset C(w)$. In Theorem 1.2 of [?], Laurence gives a formula for the centralizer of a cyclically reduced element $x \in A_\Gamma$. An easy corollary of his formula shows that if x is a product of two commuting elements, $x = x_1 x_2$, with disjoint support (i.e. $x_i \in A_{\Theta_i}$, with Θ_1 and Θ_2 disjoint, commuting set of vertices), then $C(x) = C(x_1) \cap C(x_2)$. In particular, we have $C(vg) = C(v) \cap C(g) \subset A_{J_{[v]}}$. Applying the automorphism ϕ_w^{-1} , then gives $C(v) \subset A_{J_{[v]}}$.

On the other hand, we also have $A_{[w]} \subset C(v)$, so applying ϕ_w gives $A_{[w]} \subset C(vg) \subset C(g)$. The centralizer of an element in a non-abelian free group cannot contain the entire free group, so we conclude that w must lie in V_{ab} , and it follows that $C(g) = A_{J_{[v]}} = C(w)$. Combining these we get $C(v) \subset C(w)$, or equivalently, $st(v) \subset st(w)$. Thus $[v] < [w]$ and $[v]$ is leaf-like.

In light of the discussion above, we can compose ϕ with an element of the leaf transvection group to obtain an outer automorphism $\tilde{\phi}$ such that for every $[w] \in \Gamma_0$, there is a representative $\tilde{\phi}_w$ which preserves $A_{J_{[w]}}$ and acts as the identity on $L_{[w]}$. If $[v]$ is adjacent to $[w]$ in Γ_0 , then $[w]$ is contained in $L_{[v]}$ so there is another representative $\tilde{\phi}_v$ which preserves $A_{J_{[v]}}$ and acts as the identity on $[w]$. (Here we are using the hypothesis that Γ_0 contains at least two vertices.) These two representatives differ by an inner automorphism which preserves $A_{J_{v,w}}$, so $\tilde{\phi}_w$ acts on $[w]$ as conjugation by an element $g \in N(J_{v,w}) \subset A_{J_{[w]}}$. Since $[w]$ commutes with $L_{[w]}$, we may assume that g lies in $A_{[w]}$. We conclude that the restriction of $\tilde{\phi}_w$ to $A_{J_{[w]}}$ is conjugation by an element of $A_{[w]}$, an inner automorphism. This holds for all maximal $[w]$, hence $\tilde{\phi}$ lies in K . \square

It remains to consider the special case in which Γ_0 consists of a single vertex.

LEMMA 4.3. *The following are equivalent*

- (i) Γ_0 consists of a single vertex $[v]$.
- (ii) $\Gamma = J_{[v]}$ for some $v \in V_{ab}$.
- (iii) The center of A_Γ is non-trivial.

Proof. (ii) \Rightarrow (iii) follows from Lemma ???. (iii) \Rightarrow (i) is clear since if $v \in \Gamma^\perp$, then $st(v) = \Gamma$ so $[v]$ is necessarily the unique maximal class. For (i) \Rightarrow (ii), note that for any $u \in L_{[v]}$, $[v] \subset lk(u)$. If $[v]$ is non-abelian, then $[v] \not\subset st(v)$, so $[u] \not\subset [v]$. Hence if $[v]$ is the unique vertex in Γ_0 , it must be abelian, and it follows from Lemma ?? that $\Gamma = J_{[v]}$. \square

PROPOSITION 4.4. *If Γ_0 consists of a single vertex $[v]$, then*

$$Out(A_\Gamma) = Tr \times (GL(A_{[v]}) \times Out(A_{L_{[v]}}))$$

where Tr is the free abelian group generated by the leaf transvections $t(u, w)$ with $u \in L_{[v]}$ and $w \in [v]$.

Proof. Any outer automorphism of A_Γ preserves the center $A_{[v]}$ and projects via $P_{[v]}$ to $Out(A_{L_{[v]}})$. This gives a homomorphism

$$Out(A_\Gamma) \rightarrow GL(A_{[v]}) \times Out(A_{L_{[v]}})$$

which is clearly split surjective. It is easy to check that the kernel of this homomorphism is the group generated by leaf transvections. \square

5. Virtual cohomological dimension

If Γ is discrete or is a complete graph, then A_Γ is free or free abelian and $Out(A_\Gamma)$ is known to have torsion-free subgroups of finite index, so that the *virtual cohomological dimension* (vcd) of $Out(A_\Gamma)$ is defined. In this section we prove that the same is true for arbitrary Γ , and we show furthermore that the vcd of $Out(A_\Gamma)$ is always finite.

We define the *dimension* of A_Γ to be the maximal rank of a free abelian subgroup of A_Γ . This is determined by the number of vertices in the largest complete subgraph of Γ , and we also call this the *dimension* of Γ . Thus Γ has dimension 1 if and only if A_Γ is free. If Γ has dimension $n > 1$, then links of vertices in Γ have dimension at most $n - 1$. Since the image of P lies in the product of $Out(A_{L_{[v]}})$, it is natural to try to use P together with inductive arguments to prove properties of $Out(A_\Gamma)$. Even for connected Γ , the subgraphs $L_{[v]}$ are not in general connected, so we must consider the disconnected case. In dimension 2 the links always generate free groups and the theory of $Out(F_n)$ comes into play; to deal with the general case we must also appeal to the result of Guirardel and Levitt [?] stated below.

If Γ has j components consisting of a single point and k components, $\Gamma_1, \dots, \Gamma_k$ consisting of more than one point, then A_Γ splits as a free product

$$A_\Gamma = F_j * A_1 * \dots * A_k$$

where F_j is a free group and A_i is the right-angled Artin group associated to Γ_i .

THEOREM 5.1 Guirardel-Levitt. *Suppose G is a group which decomposes as a free product $G = G_1 * \dots * G_n$ with at least one factor non-free. Assume that G_i and $G_i/Z(G_i)$ are torsion-free (respectively, have finite virtual cohomological dimension) for all i . If the outer automorphism groups $Out(G_i)$ are virtually torsion-free (respectively, have finite virtual cohomological dimension) for each factor G_i , then the same is true for $Out(G)$.*

We are now in a position to prove our theorem.

THEOREM 5.2. *For any finite simplicial graph Γ , the group $Out(A_\Gamma)$ is virtually torsion-free and has finite virtual cohomological dimension.*

Proof. We proceed by induction on the dimension of Γ . If Γ has dimension 1, then A_Γ is free and the theorem follows from [?].

Now suppose that Γ is connected and has dimension ≥ 2 . First assume Γ_0 has more than one vertex and consider the homomorphism P defined in Section ???. By induction, for every maximal $[v]$, $Out(A_{L_{[v]}})$ is virtually torsion-free and has finite vcd, so the same holds for the image of P . By Theorem ??, the kernel K_P is a finitely generated free abelian group, so in particular, it is torsion-free and has finite cohomological dimension. It now follows immediately that $Out^0(A_\Gamma)$ is virtually torsion-free and by the Serre spectral sequence, it has finite vcd. Since $Out^0(A_\Gamma)$ is finite index in $Out(A_\Gamma)$, the same holds for the larger group.

If Γ_0 consists of a unique vertex $[v]$, we use Proposition ???. By induction, $Out(A_{L[v]})$ is virtually torsion-free and has finite vcd, and the same is classically true for $Out(A_{[v]}) = GL(A_{[v]})$. Since the transvection group Tr is free abelian, the theorem follows as above.

Finally, applying Theorem ??, these results extend to n -dimensional graphs Γ with more than one component. This completes the induction. \square

In [?], the authors and J. Crisp studied the case in which Γ is connected and 2-dimensional. They obtained explicit upper and lower bounds on the vcd of $Out(A_\Gamma)$ and constructed a contractible “outer space” with a proper $Out(A_\Gamma)$ action. In a forthcoming paper with K.-U. Bux, the authors determine the exact vcd for many 2-dimensional right-angled Artin groups, in particular those whose defining graph is a tree [?].

It was also shown in [?] that for connected, 2-dimensional Γ , $Out(A_\Gamma)$ satisfies the Tits alternative. One would like to do an inductive argument as above to show that this holds for all Γ . However, in this case, we do not have the analogue of Theorem [?] to pass from the connected to the disconnected case.

Acknowledgements: The authors would like to thank Eddy Godelle for useful comments.

References

- BCV** Kai-Uwe Bux, Ruth Charney and Karen Vogtmann, *Automorphisms of tree-based RAAGS and partially symmetric automorphisms of free groups*, to appear in *Groups, Geometry and Dynamics*.
- CCV** Ruth Charney, John Crisp and Karen Vogtmann, *Automorphisms of 2-dimensional right-angled Artin groups*, *Geom. and Topology* **11** (2007), 2227–2264.
- CV86** Marc Culler and Karen Vogtmann, *Moduli of graphs and automorphisms of free groups*, *Invent. Math.* **84** (1986), no. 1, 91–119.
- DKR** Andrew Duncan, Ilya Kazachkov and Vladimir Remeslennikov, *Orthogonal systems of finite graphs*, arXiv:0707.0087
- God03** Eddy Godelle, *Parabolic subgroups of Artin groups of type FC*, *Pacific J. Math.* **208** (2003), no. 2, 243–254.
- GL07** Vincent Guirardel and Gilbert Levitt, *The Outer space of a free product*, *Proc. Lond. Math. Soc.* (3) **94** (2007), 695–714.
- Lau95** Michael R. Laurence, *A generating set for the automorphism group of a graph group*, *J. London Math. Soc.* (2) **52** (1995), no. 2, 318–334.
- Ser89** Herman Servatius, *Automorphisms of graph groups*, *J. Algebra* **126** (1989), no. 1, 34–60.

Ruth Charney
 Mathematics Department
 Brandeis University
 Waltham, MA 02454-9110
 U.S.A.

charney@brandeis.edu

Karen Vogtmann
 Mathematics Department
 Cornell University
 Ithaca, NY 14853-4201
 U.S.A.

vogtmann@math.cornell.edu